

REFERENCES

- Christiano AM, Uitto J: Molecular complexity of the cutaneous basement membrane zone. Revelations from the paradigms of epidermolysis bullosa. *Exp Derm* 5:1–11, 1996a
- Christiano AM, Uitto J: Impact of molecular genetic diagnosis on dystrophic epidermolysis bullosa. *Current Opinion Dermatol* 3:225–232, 1996b
- Christiano AM, Suga Y, Greenspan DS, Ogawa H, Uitto J: Premature termination codons on both alleles of the type VII collagen gene (COL7A1) in three brothers with recessive dystrophic epidermolysis bullosa. *J Clin Invest* 95:1328–1334, 1995
- Christiano AM, Hoffman GG, Zhang X, Xu Y, Tamai Y, Greenspan DS, Uitto J: Strategy for identification of sequence variants in COL7A1, and a novel 2-bp deletion mutation in recessive dystrophic epidermolysis bullosa. *Human Mut* 10:408–414, 1997
- Fine J-D, Bauer EA, Briggaman RA, et al.: Revised clinical and laboratory criteria for subtypes of inherited epidermolysis bullosa: a consensus report by the subcommittee on diagnosis and classification of the National Epidermolysis Bullosa Registry. *J Am Acad Dermatol* 24:119–135, 1991
- Hovnanian A, Rochat A, Bodemer C, et al.: Characterization of 18 new mutations in COL7A1 in recessive dystrophic epidermolysis bullosa provides evidence for distinct molecular mechanisms underlying defective anchoring fibril formation. *Am J Hum Genet* 61:599–610, 1997
- McGrath JA, Ishida-Yamamoto A, O'Grady A, Leigh IM, Eady RAJ: Structural variations in anchoring fibrils in dystrophic epidermolysis bullosa: correlation with type VII collagen expression. *J Invest Dermatol* 100:366–372, 1993
- Sakai LY, Keene DR, Morris NP, Burgeson RE: Type VII collagen is a major structural component of anchoring fibrils. *J Cell Biol* 103:1577–1586, 1986
- Shimizu H, McGrath JA, Christiano AM, Nishikawa T, Uitto J: Molecular basis of recessive dystrophic epidermolysis bullosa: Genotype/phenotype correlation in a case of moderate clinical severity. *J Invest Dermatol* 106:119–124, 1996
- Tamai K, Ishida-Yamamoto A, Matsuo S, et al.: Compound heterozygosity for a nonsense mutation and a splice site mutation in the type VII collagen gene (COL7A1) in recessive dystrophic epidermolysis bullosa. *Lab Invest* 76:209–217, 1997
- Uitto J, Christiano AM: Molecular genetics of the cutaneous basement membrane zone. Perspectives on epidermolysis bullosa and other blistering skin diseases. *J Clin Invest* 90:687–692, 1992
- Uitto J, Pulkkinen L, Christiano AM: The molecular basis of the dystrophic forms of epidermolysis bullosa. In: Fine J-D et al. eds. *Epidermolysis Bullosa*, Johns Hopkins University Press, Baltimore, MD, 1999, in press

Native Type I Collagen is Not a Substrate for MMP2 (Gelatinase A)

To the Editor:

A recent JID paper by Herouy *et al.* (1998) presented some interesting new data on lipodermatosclerosis. Their interpretations of the data, however, are based upon a single paper in the MMP literature (Aimes and Quigley, 1995), which claims that MMP2 (gelatinase A) is an interstitial collagenase, a claim that we have not been able to confirm. We were the first laboratory to purify, characterize, and clone human MMP2 (Seltzer *et al.*, 1981; Collier *et al.*, 1988). We found that this proteinase had absolutely no activity against helical collagen. Native helical collagen is defined by its resistance to trypsin cleavage and characteristic viscosity in solution. Subsequent work from our laboratory has shown that MMP2 is an extremely opportunistic proteinase against any collagenous sequence in which the helicity is not perfect. For example, MMP2 cleaves helical type VII collagen within the helical portion of the molecule, but in an area that has relaxed helicity (Seltzer *et al.*, 1989). When the Aimes and Quigley paper was published we confirmed our original observations using pure TIMP-free MMP2 and helical collagen shown to be trypsin resistant. Interstitial type I collagen in which the helix was relaxed enough to render it susceptible to trypsin digestion was indeed susceptible to digestion by MMP2. Intact helical collagen was not. Ohuchi *et al.* (1997) have recently reconfirmed our original findings.

Cleavage of interstitial collagen by MMP1 yields two characteristic fragments, which lose their helicity at 37°C and become soluble. Therefore, solubilization of type I collagen is not indicative of MMP2 activity, but is of MMP1 activity. Herouy *et al.* clearly show active MMP1 in their western immunoblots, indicating that cleavage of labeled type I collagen by their extracts can definitely occur.

It is important to point out that TIMP-MMP complexes have been previously described, but are separated when subjected to SDS-PAGE electrophoresis (Goldberg *et al.*, 1989; Wilhelm *et al.*, 1989). The high molecular weight band shown in the immunoblots (Fig 2) by Herouy *et al.* (1998) either are a new type of complex, or represent an artifact of antipeptide antibodies.

Jo Louise Seltzer and Arthur Z. Eisen
Division of Dermatology, Washington University School of
Medicine, St. Louis, Missouri

REFERENCES

- Aimes RT, Quigley JP: Matrix metalloproteinase-2 is an interstitial collagenase. Inhibitor-free enzyme catalyzes the cleavage of collagen fibrils and soluble native type I collagen generating the specific 3/4 and 1/4 length fragments. *J Biol Chem* 270:5872–5876, 1995
- Collier IE, Wilhelm SM, Eisen AZ *et al.*: H-ras oncogene-transformed human bronchial-epithelial cells (TBE-1) secrete a single metalloproteinase capable of degrading basement membrane collagen. *J Biol Chem* 263:6579–6587, 1988
- Goldberg GI, Marmer BL, Grant GA, Eisen AZ, Wilhelm S, He C: Human 72-kilodalton type IV collagenase forms a complex with a tissue inhibitor of metalloproteinases designated TIMP-2. *Proc Natl Acad Sci USA* 86:8207–8211, 1989
- Herouy Y, May AE, Ponschlegel G *et al.*: Lipodermatosclerosis is characterized by elevated expression and activation of matrix metalloproteinases: implications for venous ulcer formation. *J Invest Dermatol* 111:822–827, 1998
- Ohuchi E, Kazushi I, Fuji Y, Sata H, Seiki H, Okada Y: Membrane type I matrix metalloproteinase digests interstitial collagens and other extracellular matrix macromolecules. *J Biol Chem* 272:2446–2451, 1997
- Seltzer JL, Adams SA, Grant GA, Eisen AZ: Purification and properties of a gelatin-specific neutral protease from human skin. *J Biol Chem* 256:4662–4668, 1981
- Seltzer JL, Eisen AZ, Bauer EA, Burgeson RE: Cleavage of type VII collagen by interstitial collagenase and type IV collagenase (gelatinase) derived from human skin. *J Biol Chem* 264:3822–3826, 1989
- Wilhelm SM, Collier IE, Marmer BL, Eisen AZ, Grant GA, Goldberg GI: SV40-transformed human lung fibroblasts secrete a 92 kDa type IV collagenase which is identical to that secreted by normal human macrophages. *J Biol Chem* 264:17213–17221, 1989

Reply

Lipodermatosclerosis is characterized by distinct morphologic features. Prominent perivascular fibrin cuffs and deposition of hemosiderin pigments due to red blood cell extravasation are major tissue changes occurring in patients with severe chronic venous insufficiency. Perivascular cuffs are highly organized structures primarily composed of fibrin, laminin, fibronectin, tenascin and type I and III collagens (Herrick *et al.*, 1992). Fibrous scar tissue associated with fragmentation of elastic fibers and loss of papillary structures at the dermal-epidermal junction are further histologic findings. The dermal-epidermal junction is built up by an extraordinarily complex network of interconnecting proteins such as different collagen subtypes and laminin (Burgeson and Christiano, 1997). To understand the molecular basis of these morphologic alterations